



Perspectives on Perfect and Practical Play of Pig

Todd W. Neller
Gettysburg College

<http://cs.gettysburg.edu/~tneller/papers/talks/haccpig.pdf>

Sow What's This All About?



- Introduction to the Dice Game “Pig”
- Odds and Ends: Playing to Score
- Perfect Play: Playing to Win
 - “Piglet” example
 - Value Iteration
- Practical Play: How we can play well

The Game of Pig



- The first player to reach 100 points wins.
- On a turn, a player rolls a die repeatedly until:
 - the player holds, scoring the sum of the rolls (“turn total”), or
 - a 1 (“pig”) is rolled, and there is no score change.
- Example turns:
 - roll 4, roll 5, roll 2, hold → add $4 + 5 + 2 = 11$ to score
 - roll 3, roll 6, roll 6, roll 1 → score remains the same

Pig Preliminaries



- Player's decision is always to *roll/hold*
 - *Roll* – possibly increase turn total, or lose it
 - *Hold* – definitely score current turn total
 - Pig is the simplest of a class of *jeopardy dice games*; ancestor of *Pass the Pigs*
- Hold at 20 - a simple policy that maximizes expected points per turn

Playing to Score



- Simple odds argument
 - Roll until you risk more than you stand to gain.
 - “Hold at 20”
 - 1/6 of time: $-20 \rightarrow -20/6$
 - 5/6 of time: $+4$ (avg. of 2,3,4,5,6) $\rightarrow +20/6$

Hold at 20?



- Is there a situation in which you wouldn't want to hold at 20?
 - Your score: 99; you roll 2
 - Case scenario
 - you: 79 opponent: 99
 - Your turn total stands at 20

What's Wrong With Playing to Score?



- It's mathematically optimal!
- But what are we optimizing?
- Playing to score \neq Playing to win
- Optimizing score gain per turn \neq Optimizing probability of a win

Piglet



- Simpler version of Pig with a coin
- Object: First to score 10 points
- On your turn, flip until:
 - You flip tails, and score NOTHING.
 - You hold, and KEEP the # of heads.
- Even simpler: play to 2 points

Essential Information



- What is the information I need to make a fully informed decision?
 - My score
 - The opponent's score
 - My "turn total"

A Little Notation



- $P_{i,j,k}$ – probability of a win if
i = my score
j = the opponent's score
k = my turn total
- Hold: $P_{i,j,k} = 1 - P_{j,i+k,0}$
- Flip: $P_{i,j,k} = \frac{1}{2}(1 - P_{j,i,0}) + \frac{1}{2} P_{i,j,k+1}$

Assume Rationality



- To make a smart player, assume a smart opponent.
- (To make a smarter player, know your opponent.)
- $P_{i,j,k} = \max(1 - P_{j,i+k,0}, \frac{1}{2}(1 - P_{j,i,0} + P_{i,j,k+1}))$
- Probability of win based on best decisions in any state

The Whole Story



$$P_{0,0,0} = \max(1 - P_{0,0,0}, \frac{1}{2}(1 - P_{0,0,0} + P_{0,0,1}))$$

$$P_{0,0,1} = \max(1 - P_{0,1,0}, \frac{1}{2}(1 - P_{0,0,0} + P_{0,0,2}))$$

$$P_{0,1,0} = \max(1 - P_{1,0,0}, \frac{1}{2}(1 - P_{1,0,0} + P_{0,1,1}))$$

$$P_{0,1,1} = \max(1 - P_{1,1,0}, \frac{1}{2}(1 - P_{1,0,0} + P_{0,1,2}))$$

$$P_{1,0,0} = \max(1 - P_{0,1,0}, \frac{1}{2}(1 - P_{0,1,0} + P_{1,0,1}))$$

$$P_{1,1,0} = \max(1 - P_{1,1,0}, \frac{1}{2}(1 - P_{1,1,0} + P_{1,1,1}))$$

The Whole Story



$$P_{0,0,0} = \max(1 - P_{0,0,0}, \frac{1}{2}(1 - P_{0,0,0} + P_{0,0,1}))$$

$$P_{0,0,1} = \max(1 - P_{0,1,0}, \frac{1}{2}(1 - P_{0,0,0} + P_{0,0,2}))$$

$$P_{0,1,0} = \max(1 - P_{1,0,0}, \frac{1}{2}(1 - P_{1,0,0} + P_{0,1,1}))$$

$$P_{0,1,1} = \max(1 - P_{1,1,0}, \frac{1}{2}(1 - P_{1,0,0} + P_{0,1,2}))$$

$$P_{1,0,0} = \max(1 - P_{0,1,0}, \frac{1}{2}(1 - P_{0,1,0} + P_{1,0,1}))$$

$$P_{1,1,0} = \max(1 - P_{1,1,0}, \frac{1}{2}(1 - P_{1,1,0} + P_{1,1,1}))$$

These are **winning** states!

The Whole Story



$$P_{0,0,0} = \max(1 - P_{0,0,0}, \frac{1}{2}(1 - P_{0,0,0} + P_{0,0,1}))$$

$$P_{0,0,1} = \max(1 - P_{0,1,0}, \frac{1}{2}(1 - P_{0,0,0} + 1))$$

$$P_{0,1,0} = \max(1 - P_{1,0,0}, \frac{1}{2}(1 - P_{1,0,0} + P_{0,1,1}))$$

$$P_{0,1,1} = \max(1 - P_{1,1,0}, \frac{1}{2}(1 - P_{1,0,0} + 1))$$

$$P_{1,0,0} = \max(1 - P_{0,1,0}, \frac{1}{2}(1 - P_{0,1,0} + 1))$$

$$P_{1,1,0} = \max(1 - P_{1,1,0}, \frac{1}{2}(1 - P_{1,1,0} + 1))$$

Simplified...

The Whole Story



$$P_{0,0,0} = \max(1 - P_{0,0,0}, \frac{1}{2}(1 - P_{0,0,0} + P_{0,0,1}))$$

$$P_{0,0,1} = \max(1 - P_{0,1,0}, \frac{1}{2}(2 - P_{0,0,0}))$$

$$P_{0,1,0} = \max(1 - P_{1,0,0}, \frac{1}{2}(1 - P_{1,0,0} + P_{0,1,1}))$$

$$P_{0,1,1} = \max(1 - P_{1,1,0}, \frac{1}{2}(2 - P_{1,0,0}))$$

$$P_{1,0,0} = \max(1 - P_{0,1,0}, \frac{1}{2}(2 - P_{0,1,0}))$$

$$P_{1,1,0} = \max(1 - P_{1,1,0}, \frac{1}{2}(2 - P_{1,1,0}))$$

And simplified more into a **hamsome** set of equations...

How to Solve It?



$$P_{0,0,0} = \max(1 - P_{0,0,0}, \frac{1}{2}(1 - P_{0,0,0} + P_{0,0,1}))$$

$$P_{0,0,1} = \max(1 - P_{0,1,0}, \frac{1}{2}(2 - P_{0,0,0}))$$

$$P_{0,1,0} = \max(1 - P_{1,0,0}, \frac{1}{2}(1 - P_{1,0,0} + P_{0,1,1}))$$


$$P_{0,1,1} = \max(1 - P_{1,1,0}, \frac{1}{2}(2 - P_{1,0,0}))$$

$$P_{1,0,0} = \max(1 - P_{0,1,0}, \frac{1}{2}(2 - P_{0,1,0}))$$

$$P_{1,1,0} = \max(1 - P_{1,1,0}, \frac{1}{2}(2 - P_{1,1,0}))$$

$P_{0,0,0}$ depends on $P_{0,0,1}$ depends on $P_{0,1,0}$ depends on $P_{0,1,1}$
depends on $P_{1,0,0}$ depends on $P_{0,1,0}$ depends on ...

A System of Pigquations



Dependencies between non-winning states

How Bad Is It?



- The intersection of a set of bent hyperplanes in a hypercube
- In the general case, no known method (read: PhD research)
- Is there a method that works (without being guaranteed to work in general)?
 - Yes! Value Iteration!

Value Iteration



- Start out with some values (0's, 1's, random #'s)
- Do the following until the values converge (stop changing):
 - Plug the values into the RHS's
 - Recompute the LHS values
- That's easy. Let's do it!

Value Iteration



$$P_{0,0,0} = \max(1 - P_{0,0,0}, \frac{1}{2}(1 - P_{0,0,0} + P_{0,0,1}))$$

$$P_{0,0,1} = \max(1 - P_{0,1,0}, \frac{1}{2}(2 - P_{0,0,0}))$$

$$P_{0,1,0} = \max(1 - P_{1,0,0}, \frac{1}{2}(1 - P_{1,0,0} + P_{0,1,1}))$$

$$P_{0,1,1} = \max(1 - P_{1,1,0}, \frac{1}{2}(2 - P_{1,0,0}))$$

$$P_{1,0,0} = \max(1 - P_{0,1,0}, \frac{1}{2}(2 - P_{0,1,0}))$$

$$P_{1,1,0} = \max(1 - P_{1,1,0}, \frac{1}{2}(2 - P_{1,1,0}))$$

- Assume $P_{i,j,k}$ is 0 unless it's a win
- Repeat: Compute RHS's, assign to LHS's

Iteration 1



Initially, $P_{i,j,k} = 0$

$$P_{0,0,0} = \max(1 - P_{0,0,0}, \frac{1}{2}(1 - P_{0,0,0} + P_{0,0,1}))$$

$$P_{0,0,1} = \max(1 - P_{0,1,0}, \frac{1}{2}(2 - P_{0,0,0}))$$

$$P_{0,1,0} = \max(1 - P_{1,0,0}, \frac{1}{2}(1 - P_{1,0,0} + P_{0,1,1}))$$

$$P_{0,1,1} = \max(1 - P_{1,1,0}, \frac{1}{2}(2 - P_{1,0,0}))$$

$$P_{1,0,0} = \max(1 - P_{0,1,0}, \frac{1}{2}(2 - P_{0,1,0}))$$

$$P_{1,1,0} = \max(1 - P_{1,1,0}, \frac{1}{2}(2 - P_{1,1,0}))$$

Iteration 1



Initially, $P_{i,j,k} = 0$

$$P_{0,0,0} = \max(1 - 0, \frac{1}{2}(1 - 0 + 0))$$

$$P_{0,0,1} = \max(1 - 0, \frac{1}{2}(2 - 0))$$

$$P_{0,1,0} = \max(1 - 0, \frac{1}{2}(1 - 0 + 0))$$

$$P_{0,1,1} = \max(1 - 0, \frac{1}{2}(2 - 0))$$

$$P_{1,0,0} = \max(1 - 0, \frac{1}{2}(2 - 0))$$

$$P_{1,1,0} = \max(1 - 0, \frac{1}{2}(2 - 0))$$

Iteration 1



Initially, $P_{i,j,k} = 0$

$$P_{0,0,0} = \max(1, \frac{1}{2}) = 1$$

$$P_{0,0,1} = \max(1, 1) = 1$$

$$P_{0,1,0} = \max(1, \frac{1}{2}) = 1$$

$$P_{0,1,1} = \max(1, 1) = 1$$

$$P_{1,0,0} = \max(1, 1) = 1$$

$$P_{1,1,0} = \max(1, 1) = 1$$

Iteration 2



Next, $P_{i,j,k} = 1$

$$P_{0,0,0} = \max(1 - P_{0,0,0}, \frac{1}{2}(1 - P_{0,0,0} + P_{0,0,1}))$$

$$P_{0,0,1} = \max(1 - P_{0,1,0}, \frac{1}{2}(2 - P_{0,0,0}))$$

$$P_{0,1,0} = \max(1 - P_{1,0,0}, \frac{1}{2}(1 - P_{1,0,0} + P_{0,1,1}))$$

$$P_{0,1,1} = \max(1 - P_{1,1,0}, \frac{1}{2}(2 - P_{1,0,0}))$$

$$P_{1,0,0} = \max(1 - P_{0,1,0}, \frac{1}{2}(2 - P_{0,1,0}))$$

$$P_{1,1,0} = \max(1 - P_{1,1,0}, \frac{1}{2}(2 - P_{1,1,0}))$$

Iteration 2



Next, $P_{i,j,k} = 1$

$$P_{0,0,0} = \max(1 - 1, \frac{1}{2}(1 - 1 + 1))$$

$$P_{0,0,1} = \max(1 - 1, \frac{1}{2}(2 - 1))$$

$$P_{0,1,0} = \max(1 - 1, \frac{1}{2}(1 - 1 + 1))$$

$$P_{0,1,1} = \max(1 - 1, \frac{1}{2}(2 - 1))$$

$$P_{1,0,0} = \max(1 - 1, \frac{1}{2}(2 - 1))$$

$$P_{1,1,0} = \max(1 - 1, \frac{1}{2}(2 - 1))$$

Iteration 2



Next, $P_{i,j,k} = 1$

$$P_{0,0,0} = \max(0, \frac{1}{2}) = \frac{1}{2}$$

$$P_{0,0,1} = \max(0, \frac{1}{2}) = \frac{1}{2}$$

$$P_{0,1,0} = \max(0, \frac{1}{2}) = \frac{1}{2}$$

$$P_{0,1,1} = \max(0, \frac{1}{2}) = \frac{1}{2}$$

$$P_{1,0,0} = \max(0, \frac{1}{2}) = \frac{1}{2}$$

$$P_{1,1,0} = \max(0, \frac{1}{2}) = \frac{1}{2}$$

Iteration 3



Next, $P_{i,j,k} = \frac{1}{2}$

$$P_{0,0,0} = \max(1 - P_{0,0,0}, \frac{1}{2}(1 - P_{0,0,0} + P_{0,0,1}))$$

$$P_{0,0,1} = \max(1 - P_{0,1,0}, \frac{1}{2}(2 - P_{0,0,0}))$$

$$P_{0,1,0} = \max(1 - P_{1,0,0}, \frac{1}{2}(1 - P_{1,0,0} + P_{0,1,1}))$$

$$P_{0,1,1} = \max(1 - P_{1,1,0}, \frac{1}{2}(2 - P_{1,0,0}))$$

$$P_{1,0,0} = \max(1 - P_{0,1,0}, \frac{1}{2}(2 - P_{0,1,0}))$$

$$P_{1,1,0} = \max(1 - P_{1,1,0}, \frac{1}{2}(2 - P_{1,1,0}))$$

Iteration 3



Next, $P_{i,j,k} = \frac{1}{2}$

$$P_{0,0,0} = \max(1 - \frac{1}{2}, \frac{1}{2}(1 - \frac{1}{2} + \frac{1}{2}))$$

$$P_{0,0,1} = \max(1 - \frac{1}{2}, \frac{1}{2}(2 - \frac{1}{2}))$$

$$P_{0,1,0} = \max(1 - \frac{1}{2}, \frac{1}{2}(1 - \frac{1}{2} + \frac{1}{2}))$$

$$P_{0,1,1} = \max(1 - \frac{1}{2}, \frac{1}{2}(2 - \frac{1}{2}))$$

$$P_{1,0,0} = \max(1 - \frac{1}{2}, \frac{1}{2}(2 - \frac{1}{2}))$$

$$P_{1,1,0} = \max(1 - \frac{1}{2}, \frac{1}{2}(2 - \frac{1}{2}))$$

Iteration 3



Next, $P_{i,j,k} = \frac{1}{2}$

$$P_{0,0,0} = \max(\frac{1}{2}, \frac{1}{2}) = \frac{1}{2}$$

$$P_{0,0,1} = \max(\frac{1}{2}, \frac{3}{4}) = \frac{3}{4}$$

$$P_{0,1,0} = \max(\frac{1}{2}, \frac{1}{2}) = \frac{1}{2}$$

$$P_{0,1,1} = \max(\frac{1}{2}, \frac{3}{4}) = \frac{3}{4}$$

$$P_{1,0,0} = \max(\frac{1}{2}, \frac{3}{4}) = \frac{3}{4}$$

$$P_{1,1,0} = \max(\frac{1}{2}, \frac{3}{4}) = \frac{3}{4}$$

This continues until values converge...

But That's GRUNT Work!



- So have a computer do it, slacker!
- Not difficult – end of CS1 level
- Fast! Don't blink – you'll miss it
- Optimal play:
 - Compute the probabilities
 - Determine flip/hold from RHS max's
 - (For our equations, always FLIP)

Piglet Solved




- Game to 10
- Play to Score: "Hold at 1"
- Play to Win:

Opponent

	0	1	2	3	4	5	6	7	8	9
0	2	2	2	2	2	2	2	2	3	10
1	1	2	2	2	2	2	2	3	3	9
2	1	1	2	2	2	2	2	2	3	8
3	1	1	1	2	2	2	2	2	3	7
4	1	1	1	1	2	2	2	2	2	6
5	1	1	1	1	1	2	2	2	2	5
6	1	1	1	1	1	1	2	2	2	4
7	1	1	1	1	1	1	1	2	3	3
8	1	1	1	1	1	1	1	2	2	2
9	1	1	1	1	1	1	1	1	1	1

You



Pig Probabilities



- Just like Piglet, but more possible outcomes


- $P_{i,j,k} =$
 $\max(1 - P_{j,i+k,0},$
 $1/6((1 - P_{j,i,0}) + P_{i,j,k+2} + P_{i,j,k+3}$
 $+ P_{i,j,k+4} + P_{i,j,k+5} + P_{i,j,k+6}))$

Solving Pig




- 505,000 such equations
- Same simple solution method (value iteration)
- Potential Speedup: Solve groups of interdependent probabilities from game end backward
- So what does optimal play look like?


Pig Sow-lution



Pig Sow-lution




Probability Contours



Practical Play of Pig



- Whoa! That's some funky alien landscape on that optimal policy!
- (scratches head) So I'm supposed to memorize that?
- Computing optimal play of Pig didn't make *me* play optimally.
- How does one come up with practical policies for unaided human play?



Approximating Optimality



- KISS Principle: “Keep It Simple and Stupid.”
- Often, much learning benefit comes from attention to few, simple features.
- Observe the optimal policy and look for significant features of the roll/hold boundary.
- Try, try again.
- How does one evaluate simple policy ideas?

Evaluating Policies



- Given two policies (yours and optimal):
 - set up a system of equations describing play,
 - compute the probability of your winning going first/second, and
 - average the win probabilities

Policy Comparison



Algorithm 1 Policy Comparison

For each $(i, j, k) \in \mathcal{S}$, initialize $P_{i,j,k}^A$ and $P_{i,j,k}^B$ arbitrarily.

Repeat

$\Delta \leftarrow 0$

For each $(i, j, k) \in \mathcal{S}$,

$$p1 \leftarrow \begin{cases} \frac{1}{6} \left[(1 - P_{j,i,0}^B) + \sum_{r \in [2,6]} P_{i,j,k+r}^A \right], & \text{if } \text{Roll}_{i,j,k}^A; \\ 1 - P_{j,i+k,0}^B, & \text{otherwise.} \end{cases}$$

$$p2 \leftarrow \begin{cases} \frac{1}{6} \left[(1 - P_{j,i,0}^A) + \sum_{r \in [2,6]} P_{i,j,k+r}^B \right], & \text{if } \text{Roll}_{i,j,k}^B; \\ 1 - P_{j,i+k,0}^A, & \text{otherwise.} \end{cases}$$

$$\Delta \leftarrow \max \{ \Delta, |p1 - P_{i,j,k}^A|, |p2 - P_{i,j,k}^B| \}$$

$$P_{i,j,k}^A \leftarrow p1$$

$$P_{i,j,k}^B \leftarrow p2$$

until $\Delta < \epsilon$

return $[P_{0,0,0}^A + (1 - P_{0,0,0}^B)] / 2$

Hold at n (or goal)




Figure 2. Probability of an optimal player winning against a player using the "hold at n " policy

$n = 25 \rightarrow$ optimal advantage = 4.2%; $n = 20 \rightarrow$ optimal advantage = 8.0%

t Scoring Turns



■ Hold value:
$$h(i, t_s) = \left\lfloor \frac{100 - i}{t - t_s} \right\rfloor$$




Figure 4. The probability of an optimal player winning against a player using the "t scoring turns" policy for different values of t.

$t = 4 \rightarrow$ optimal advantage = 3.3%

Score Base, Keep Pace, End Race



- Roll if:
 - $k < b$ (you must score at least base value b),
 - $i + k < j - p$ (you must get within p of j), or
 - either $i \geq 100 - e$ or else $j \geq 100 - e$ (you roll to win when someone is within e of the goal).
- Optimizing parameters, $b = 19$, $p = 14$, and $e = 31 \rightarrow$ optimal advantage = 1.9%

Keep Pace and End Race



- Roll if:
 - either $i \geq 100 - e$ or else $j \geq 100 - e$, or
 - $k < c + (j - i)/d$.
- Optimizing $c = 21$, $d = 8$, $e = 29$, and rounding division for hold value \rightarrow optimal advantage = **0.922%**
- So, if either player's score is 71 or higher, roll for the goal. Otherwise, subtract your score from your opponent's and let m be the closest multiple of 8. (Choose the greater multiple if halfway between multiples.) Then hold at $21 + m/8$.

Summary



- What we've learned:
 - Playing to score is not necessarily playing to win.
 - Simple rules do not imply simple perfect play.
 - Making a guess at a solution and iteratively improving that guess can be a useful method.
 - Similar iterative techniques can help us capture the simple essence of good play.
 - The computer is an exciting power tool for the mind!

Related Resources



The Game of Pig page:

<http://cs.gettysburg.edu/projects/pig>



Pig CS teaching resources:

<http://cs.gettysburg.edu/~tneller/resources/pig>